



One Health approach in “Do It Yourself” stores to tackle fungal contamination

Marta Dias^{1,2}; Pedro Pena^{1,2}; Bianca Gomes^{2,3}; Renata Cervantes^{1,2}; Susana Viegas^{1,2}, Carla Viegas^{1,2}

1 NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal

2 H&TRC – Health & Technology Research Center, ESTeSL – Escola Superior de Tecnologia e Saúde, Instituto Politécnico de Lisboa; carla.viegas@estesl.ipl.pt

3 CE3C—Center for Ecology, Evolution and Environmental Change, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisbon, Portugal

Introduction



➤ The DIY (Do-It-Yourself) market refers to the **products used in home renovation and repair projects (1)**

➤ The target market for these items is those **who want to conduct their home renovations (1)**



➤ SARS-CoV-2 pandemic and issues with the global supply **chain didn't stop the sector from growing (2)**

➤ The sector growth resulted from **COVID-19 limitations** that kept many people confined to their houses

Introduction





➤ The **sheer number of customers and the items' diverse origins** provide a distinct environment (3-5)

➤ It is crucial to consider the **various exposure scenarios** when focusing on microbial contamination




➤ This study aimed to assess the **exposure of woodworking employees and customers to fungal load**

Materials and Methods




Results - MEA



- **I/O:** higher indoor load compared to the outdoor in all sampling sites in two stores
- **WHO (150 CFU.m⁻³):** at least one sampling site surpasses this limit in most stores
- **OSHA (1.0 x 10¹-1.0 x 10⁴ CFU.m⁻³):** all the sampling sites from all the stores are within the limit

Results – DG18

- **I/O:** higher indoor load compared to the outdoor in five stores in all sampling sites
- **WHO (150 CFU.m⁻³):** at least one sampling site surpasses this limit in most stores
- **OSHA (1.0 x 10¹-1.0 x 10⁴ CFU.m⁻³):** all the sampling sites from all the stores are within the limit



Results

Regarding the customers' exposure and applying the Portuguese Indoor Air Quality (IAQ) legal framework (Portaria n.º 138-G/2021, de 1 de julho):


MEA

- I/O ratio complied in 5 out of 13 stores (38,5%)
- in one of the remaining 8, one toxigenic species with a quantitative cut-off above the legal frame was identified

DG18

- I/O did not comply in any store

Additionally: in 6 out of the 13 stores (46,2%), three toxigenic species with a quantitative cut-off above the legal frame were found



Discussion

- There are no occupational **exposure limits for microbial agents** (6)
- **Different results on MEA and DG18**, highlight the potential of a selective media for fungal load (7)
- **I/O ratio < 1** is used as guidance and suggests an acceptable indoor environment (8)
- I/O ratio results suggest the **need to implement measures** aiming risk control (8), like in other occupational environments (9)



Discussion



- Wood shavings have been frequently **used as animals' bedding** by the poultry industry (10)
- The bedding material is one of the **main causes of indoor air microbial contamination** in poultry (10)
- Increase on animal and occupational research in poultries, **reinforcing the One Health strategy** (6,10)

Determine priority areas for intervention to reduce microbiological exposure in DIY stores

Main Findings

- Health concerns regarding workers' and customers' safety **support the need to prevent exposure**
- Selective culture media allow us a **more detailed analysis** preventing the overgrowth of some species
- Wood dust and shavings from DIY stores **may be used in animal bedding and agriculture**
- The use of wood shavings in other environments **highlight the need for a One Health approach**

The novelty of this data is to allow a prioritized intervention by Occupational Health Services aiming an accurate exposure assessment and allowing a more detailed risk assessment and management





Escola Nacional
de Saúde Pública
UNIVERSIDADE NOVA DE LISBOA

CHRC
COMPREHENSIVE HEALTH
RESEARCH CENTRE



ESCOLA SUPERIOR DE
TECNOLOGIA DA SAÚDE
DE LISBOA
INSTITUTO POLITÉCNICO DE LISBOA

H&TRC
Health and Technology Research Center

Thank you for your attention!

Marta Dias

marta.dias@estesl.ipl.pt

msf.dias@ensp.unl.pt

Acknowledgments:

This research was funded by national funds through the FCT—Fundação para a Ciência e Tecnologia, I.P., within the scope of the PhD Grant UI/BD/151431/2021. H&TRC authors gratefully acknowledge the FCT/MCTES national support through the 2023.01366.BD; UIDB/05608/2020; UIDP/05608/2020 and PL/2022/InChildhealth/BI/12M. The Instituto Politécnico de Lisboa, national support through IPL/IDI&CA2023/FoodAllEU; IPL/IDI&CA2023/ASPRisk and IPL/IDI&CA2023/ARAFSawmil

References

1. Statista [Internet]. [citado 4 de janeiro de 2024]. DIY & Hardware Store - Worldwide | Statista Market Forecast. Disponível em: <https://www.statista.com/outlook/cmo/diy-hardware-store/worldwide>
2. Statista [Internet]. [citado 4 de janeiro de 2024]. DIY & Hardware Store - Portugal | Statista Market Forecast. Disponível em: <https://www.statista.com/outlook/cmo/diy-hardware-store/portugal>
3. Hoisington A, Maestre JP, Kinney KA, Siegel JA. Characterizing the bacterial communities in retail stores in the United States. *Indoor Air*. dezembro de 2016;26(6):857–68.
4. Frankel M, Bekö G, Timm M, Gustavsen S, Hansen EW, Madsen AM. Seasonal variations of indoor microbial exposures and their relation to temperature, relative humidity, and air exchange rate. *Appl Environ Microbiol*. dezembro de 2012;78(23):8289–97.
5. Nasir ZA, Colbeck I. Assessment of Bacterial and Fungal Aerosol in Different Residential Settings. *Water Air Soil Pollut*. setembro de 2010;211(1–4):367–77.
6. Dias, M., Pena, P., Gomes, B., Cervantes, R., Viegas, S., & Viegas, C. (2024). Snapshot Assessment to Fungal Load in Wood Departments from “Do It Yourself” (DIY) Stores. Em P. M. Arezes, R. B. Melo, P. Carneiro, J. Castelo Branco, A. Colim, N. Costa, S. Costa, J. Duarte, J. C. Guedes, G. Perestrelo, & J. S. Baptista (Eds.), *Occupational and Environmental Safety and Health V* (pp. 389–398). Springer Nature Switzerland. https://doi.org/10.1007/978-3-031-38277-2_31
7. Viegas, C. Sampling Methods for an Accurate Mycobiota Occupational Exposure Assessment: Overview of Several Ongoing Projects; Taylor & Francis: Abingdon, UK, 2018; pp. 7–11.
8. Cox, J., Mbareche, H., Lindsley, W. G., & Duchaine, C. (2020). Field sampling of indoor bioaerosols. *Aerosol Science and Technology*, 54(5), 572–584. <https://doi.org/10.1080/02786826.2019.1688759>
9. Viegas C, Gomes B, Dias M, Carolino E, Aranha Caetano L. *Aspergillus Section Fumigati* in Firefighter Headquarters. *Microorganisms*. 7 de outubro de 2021;9(10):2112.
10. Gomes B, Dias M, Cervantes R, Pena P, Santos J, Vasconcelos Pinto M, et al. One Health Approach to Tackle Microbial Contamination on Poultry—A Systematic Review. *Toxics*. abril de 2023;11(4):374.